

**YESHIVA UNIVERSITY
GRADUATE PROGRAM IN MATHEMATICAL SCIENCES
TOPICS FOR THE PHD QUALIFYING EXAMINATION**

The qualifying examination in mathematical sciences covers three areas:

- (I) Real Analysis
- (II) Complex Analysis
- (III) Research Area

For the first two areas, a list of topics is below. Also, a list of sample exercises for the two areas is provided. The actual exercises asked on the exam will be different from the sample exercises; being able to solve the sample exercises is not sufficient for the exam preparation.

The third exam area pertains to the research subject that the student intends

7. Riemann-Lebesgue Theorem (outline). A function $f : [a; b] \rightarrow \mathbb{R}$ is Riemann integrable if and only if it is bounded and its set of discontinuity points is of measure zero.
8. Sequences of functions (uniform convergence, properties, equi-continuity for a family of functions, Ascoli-Arzelà's theorem).

Complex Analysis:

1. If f is complex differentiable at z then the Cauchy-Riemann equations are satisfied at z .
2. If the partial derivatives of u and v exist and are continuous at $(x; y)$ and the Cauchy-Riemann equations are satisfied then $f(z) = u(x; y) + iv(x; y)$ is complex differentiable at $z = x + iy$.
3. If $f'(z) = 0$ in a region D then f is constant on D .
4. If $|f(z)| < M$ on a curve C then $\int_C f(z) dz < ML$ where L is the length of the curve.
5. The following statements are equivalent:
 - (i) f has an antiderivative F ;
 - (ii) $\int_{z_1}^{z_2} f(z) dz = F(z_2) - F(z_1)$;
 - (iii) If C is a closed curve then $\int_C f(z) dz = 0$.
6. Cauchy-Goursat Theorem (outline). If f is analytic on and inside a simple closed curve C then $\int_C f(z) dz = 0$.
7. If f is analytic in the region between closed curves C_2 and C_1 with C_1 inside C_2 then

$$\int_{C_1} f(z) dz = \int_{C_2} f(z) dz:$$
8. The Cauchy Integral Formula.
9. A bounded entire function is constant.
10. If f is analytic on annulus, it equals its Laurent series (outline).
11. Cauchy Residue Theorem.

Sample Exercises:

1. Let

$$f(x) = \begin{cases} \sin \frac{1}{x} & ; \text{ for } x \neq 0; \\ & ; \text{ for } x = 0, \end{cases}$$

where $x \in [-1; 1]$.

(a) Is f continuous?

(b) Does f have the intermediate value property?

2. Let

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x} & ; \text{ for } x \neq 0; \\ 0 & ; \text{ for } x = 0, \end{cases}$$

(a) Show that f is differentiable everywhere.

(b) Is f' continuous?

3. Given that f is a quadratic polynomial

$$f(x) = Kx^2 +$$

6. On uniform convergence.

(a) Show that if a sequence $\{f_n\}$ of continuous functions converges uniformly on a domain $D \subset \mathbb{R}$ to a function f , then the limit f is also continuous on D .

(b) Let $\{f_n\}$ be a sequence of continuously differentiable functions such that $\{f_n\}$ and $\{f_n'\}$ converge uniformly on a domain D to the limiting functions f and g , respectively. Show that for every x in the interior of D ,

$$g(x) = \lim_{n \rightarrow \infty} f_n'(x) = \left(\lim_{n \rightarrow \infty} f_n(x) \right)' = f'(x).$$

7. State Ascoli-Arzelà's Theorem and outline its proof.

8. The sequence of continuous functions $\{f_n\} : [0; 2\pi] \rightarrow \mathbb{R}$, $n \in \mathbb{N}$ with f_n given by $f_n(x) = \sin(nx)$ is uniformly bounded, but not equicontinuous. Give an intuitive reason why such a sequence is not equicontinuous, then give a rigorous proof.

9. Compute the following integral limit

$$\lim_{n \rightarrow \infty} \int_{-\infty}^{\infty} \frac{n \sin(x/n)}{x(x^2 + 1)} dx$$

10. Consider the function

$$f(z) = \begin{cases} \frac{z^2}{z} & \text{if } z \neq 0; \\ 0 & \text{if } z = 0; \end{cases}$$

Is this function differentiable at $z = 0$? Is it continuous at




Figure 1: Contour C

15. Show that the only conformal maps from the complex plane onto itself are the non-constant linear maps, i.e. maps of the form $f(z) = az + b$, $a \neq 0$.
16. Let f be a *doubly periodic function*, that is, there are two complex numbers w_1, w_2 with $w_1, w_2 \notin \mathbb{R}$ so that for any $z \in \mathbb{C}$, $f(z) = f(z + w_1) = f(z + w_2)$. Let us also assume that f is meromorphic.
 - (a) Show that if f is an entire function, then it has to be constant.
 - (b) Let γ be the boundary of the parallelogram with vertices $0, w_1, w_2, w_1 + w_2$, oriented counterclockwise. Show that if f is analytic on γ , then $\int_{\gamma} f(z) dz = 0$.
 - (c) Assuming that f is analytic on γ and has exactly one singularity inside γ , show that the residue at this singularity is necessarily zero.

Bibliography:

1. Charles C. Pugh. Real Mathematical Analysis. Springer.
2. Robert G. Bartle, Donald R. Sherbert. Introduction to Real Analysis Fourth Edition. John Wiley and Sons.
3. Walter Rudin. Principles of Mathematical Analysis. McGraw-Hill, Inc.
4. H.L. Royden. Real Analysis. Pearson Custom Publishing.
5. James Brown and Ruel Churchill. Complex Variables and Applications. McGraw-Hill, Inc.
6. Walter Rudin. Real and Complex Analysis. McGraw-Hill, Inc.